



EURL POPs Priority List of PCN congeners

Version 1.1

05 June 2023

Background and Rationale for the prioritisation of specific
PCN congeners in food



Document history

Action	Date	Version
Publication of Priority List of PCN congeners on EURL POPs website	31 March 2023	V 1.0
Publication of Priority List of PCN congeners on EURL POPs website	05 June 2023	V 1.1

Citation

This guidance document is made publicly available on the EURL POPs website (www.eurlpops.eu). The recommended citation for this report is as follows:

EURL for halogenated POPs in feed and food (2023): EURL POPs Priority List of PCN congeners - Background and Rationale for the prioritisation of specific PCN congeners in food, version 1.1 of 05 June 2023. Available online under <https://eurl-pops.eu/news/priority-list-pcns/priority-list-pcns>

Authorship

This priority list on PCN congener selection was prepared and discussed by members of the Core Working Group “Brominated Contaminants and Polychlorinated Naphthalenes” (CWG BCons & PCNs) of the network of the European Union Reference Laboratory (EURL) and the National Reference Laboratories (NRLs) for halogenated POPs in Feed and Food and invited experts.

Editors

Alwyn Fernandes (Independent expert, UK)

Alexander Schächtele (EURL, Germany)

Theresa Zwickel (EURL, Germany)

Members of the Working Group for Brominated Contaminants	
Thorsten Bernsmann	Chemisches und Veterinäruntersuchungsamt Münsterland-Emscher-Lippe, Münster, Germany
Tatiana Cantillana	Swedish food Agency, Sweden
Alwyn Fernandes (Chairman)	Independent expert, UK
Ali, Aasim	The Institute of Marine Research (Norwegian NRL for Halogenated POPs in feed and food), Bergen, Norway.
Leondios Leondiadis	National Centre for Scientific Research (NCSR) “Demokritos”, Athens, Greece
Philippe Marchand	Laboratoire d’Etude des Résidus et Contaminants dans les Aliments (LABERCA), France
Sebastian Maszewski	Państwowy Instytut Weterynaryjny - Państwowy Instytut Badawczy, Puławy, Poland
Panu Rantakokko	National Institute for Health and Welfare, Helsinki, Finland
Alexander Schächtele	European Union Reference Laboratory (EURL) for Halogenated POPs, Freiburg, Germany
Georges Scholl	Centre de Recherche Analytique et Technologique (CART), Belgium
Anaïs Vénisseau	Laboratoire d’Etude des Résidus et Contaminants dans les Aliments (LABERCA), France
Dzintars Zacs	Institute of Food Safety, Animal Health and Environment “BIOR”, Latvia
Theresa Zwickel	European Union Reference Laboratory (EURL) for Halogenated POPs, Freiburg, Germany

1. Description and Scope

This document summarizes background information and syntheses of supporting information on the outcome of the discussion of the EURL CWG “BCons & PCNs” meeting held on November 22, 2022. The aim was to provide a priority list of PCN congeners to assist laboratories in method development and also to support future assessments with relevant as well as harmonised data. The guidance in this document is intended for laboratories involved in the official control of contaminants in food.

2. Background

Polychlorinated naphthalenes (PCNs) are a group of 75 individual congeners that are classified as persistent organic pollutants (POPs) under the Stockholm Convention [1]. They are confirmed contaminants in foodstuffs and also occur in human tissues. At high levels of human and animal exposure, PCNs cause serious and sometimes fatal liver damage as well as dermal, neurological and endocrine effects [2,3]. Following awareness of these adverse health effects, PCN global production (estimated at ~ 1.5 m tonnes [2,4]) was phased out around three to four decades ago, but as stable chemicals, the legacy of this production endures to the present time. Current human exposure has been reported to occur mainly through dietary intake and in order to evaluate this risk, surveillance of the food supply for PCN content would be a necessary first step. The 75 congeners show varying degrees of toxicity [2,5] so prioritising the more toxicologically significant and more abundantly occurring congeners would allow a pragmatic approach to the monitoring of food.

3. Synthesis of the supporting information

There are a number of independent studies from different global locations that have reported on the occurrence of PCNs in food over the last decade [2,6-9]. These studies have reported data for a varying number of PCN congeners due to the limited availability of reliable analytical standards. Nevertheless, most studies were able to include the more toxicologically significant congeners. More analytical standards have since become available. However, there is no structured monitoring in any country that would allow a current estimate of the background content in food or the visualisation of a trend in occurrence levels, which is essential for reliable risk assessment. In order to support future assessments with relevant as well as harmonised data, the EURL CWG for BCons and PCNs has focussed on developing a priority list of congeners for the monitoring of PCNs in food. This process considered the available occurrence data as well as recent reports on the toxic effects of PCNs [2,5,10]. These describe a range of effects, some occurring at higher levels of PCN dosing that would be expected from occupational exposure. However, as most current exposure is expected to result from chronic low-level dietary intake, it may be more appropriate to consider health effects that correspond to these levels. Thus, the CWG’s evaluation focussed on effects arising from interaction with the aryl hydrocarbon receptor (dioxin-like effects). Based on the available congener relative potency (REP) data (derived mostly from in vitro studies and one in silico study) [2,5,11] and the occurrence distribution of PCNs in foods and in human milk, it was possible to prioritise a set of 19 PCN congeners for food monitoring studies (Table 1). If the concentrations of these congeners were converted to toxic equivalents (TEQs) using the available REP data [2], the summed TEQ of the 19 congeners would represent >98% of the toxic equivalence of PCNs in



the foods (>99% in human milk). It was not possible to establish a similar set of priority congeners for the monitoring of animal feeds – the PCN database for these materials is very limited and also shows variability in occurrence, an aspect that was not prominent in the food data. However, the CWG will investigate additional PCN congeners (particularly some tri- to penta-chlorinated congeners) that may be more relevant to feed monitoring. Currently, it is very clear that substantially more data on the PCN content of animal feeds is required before a similar priority list for feed can be proposed.

The PCN congener patterns from the available occurrence data on food and human tissues [8,9,12] indicate that the dominant source is likely to be associated with the legacy of PCN manufacture and widespread use in Europe and North America. However, some regions (e.g., some parts of East Asia) where PCNs have not been manufactured or used to any significant extent may show different patterns of occurrence [11], such as those associated with combustion sources.

Table 1: PCN congener priority list for monitoring of food of the network of EURL and NRLs for halogenated POPs in feed and food

Chlorination degree	Name	Congener No.
Tetra (Te)	1,2,3,5 – TeCN	28
	1,2,5,6 – TeCN	36
	1,3,5,7 – TeCN	42
	2,3,6,7 – TeCN	48
Penta (Pe)	1,2,3,5,7 – PeCN	52
	1,2,3,5,8 – PeCN	53
	1,2,4,5,8 – PeCN	59
	1,2,4,6,7 – PeCN	60
Hexa (Hx)	1,2,3,4,5,6 – HxCN	63
	1,2,3,4,5,7 – HxCN	64
	1,2,3,4,6,7 – HxCN	66
	1,2,3,5,6,7 – HxCN	67
	1,2,3,5,6,8 – HxCN	68
	1,2,3,5,7,8 – HxCN	69
	1,2,3,6,7,8 – HxCN	70
	1,2,4,5,6,8 – HxCN	71
1,2,4,5,7,8 – HxCN	72	
Hepta (Hp)	1,2,3,4,5,6,7 – HpCN	73
	1,2,3,4,5,6,8 – HpCN	74
Octa (O)	1,2,3,4,5,6,7,8 – OCN	75

4. REFERENCES

1. UNEP, 2019. The New POPs under the Stockholm Convention. <http://www.pops.int/>
2. Fernandes, A.R., Kilanowicz, A., Stragierowicz, J., Klimczak, M. and Falandysz, J., 2022. The toxicological profile of polychlorinated naphthalenes (PCNs). *Sci. Tot. Environ.* 837, 155764.
3. Ward, E., Ruder, A., Suruda, A., Smith, A., Fessler-Flesch, C., Zahm, S. 1996. Acute and chronic liver toxicity resulting from exposure to chlorinated naphthalenes at a cable manufacturing plant during World War II. *Am. J. Ind. Med.* 30(2), 225-233.
4. Falandysz, J., 2003. Chloronaphthalenes as food-chain contaminants: a review. *Food Addit. Contam.* 20, 995–1014.
5. Suzuki, G., Michinaka, C., Matsukami, H., Noma, Y. Kajiwara, N., 2020. Validity of using a relative potency factor approach for the risk management of dioxin-like polychlorinated naphthalenes. *Chemosphere*, 244, 125448.
6. Fernandes, A., Mortimer, D., Gem, M., Smith, F., Rose, M., Panton, S., Carr, M. 2010. Polychlorinated naphthalenes (PCNs): congener specific analysis, occurrence in food and dietary exposure in the UK. *Environ. Sci. Technol.* 44, 3533–3538.
7. Fernandes, A., Tlustos, C., Rose, M., Smith, F., Carr, M., Panton S. 2011. Polychlorinated naphthalenes (PCNs) in Irish foods: occurrence and human exposure. *Chemosphere* 85, 322–328.
8. Fernandes, A., Rose, M., Falandysz J. 2017. PCNs in food and humans. *Environ. Int.* 104, 1–13.
9. Zacs, D., Perkons, I., Pasecnaja, E., Bartkevics, V. 2021. Polychlorinated naphthalenes (PCNs) in food products in Latvia: congener specific analysis, occurrence, and dietary exposure of the general population. *Chemosphere* 264, 128460.
10. Falandysz, J., Fernandes, A, Gregoraszczyk, E, Rose, M. 2014. The toxicological effects of halogenated naphthalenes - a review of aryl hydrocarbon receptor mediated (dioxin-like) relative potency factors. *J. Environ. Sci. Health. C* 32, 239–272.
11. Li, C., Zhang, L., Yang, Q., Wu, Y., Zheng, M., et al., 2022. Comprehensive Evaluation of Dietary Exposure and Health Risk of Polychlorinated Naphthalenes. *Environ. Sci. Technol.* 56(9), 5520-5529
12. Tschiggfrei et al., 2023 Legacy and emerging persistent organic pollutants in human milk. *Handbook Environ. Chem.* (in press).